



1 Numbers and Sets

1.1 Sets

Set: collection of distinct objects which are called elements (numbers, people, letters of alphabet)

Two ways of defining sets:

- list each member of the set (e.g. $\{4,2,15,6\}$, $\{\text{red, blue, white}\}$, ...)

The order in which the elements of a set are listed is irrelevant, as are any repetitions in the list. For example,

$$\{6, 11\} = \{11, 6\} = \{11, 11, 6, 11\}$$

are equivalent, because the specification means merely that each of the elements listed is a member of the set.

- rule (e.g. $A = \text{set of even numbers}$, $B = \{n^2, n \in \mathbb{N}, 0 \leq n \leq 5\}$, ...)

Membership: some elements belong to a set and some do not


- $4 \in A$, $15 \in \{4, 2, 15, 6\}$, $16 \in B$ (read " \in " as "belongs to")
- $5 \notin A$, $5 \notin B$, $\text{green} \notin \{\text{red, blue, white}\}$ (read " \notin " as "does not belong to")

Cardinality: the number of members of a set

- $|A| = \infty$ (The set A has infinitely many elements)
- $|B| = 6$ (The set B has six elements)
- $|C| = 0$, where $C = \{\text{three sided squares}\}$ (The set C is an empty set)

Subsets:

- $A \subseteq B$ if every member of A is in B as well (A is subset of B)
- if $A \subseteq B$ but $A \neq B$, then A is a proper subset of B , $A \subset B$
- $\{1, 2\} \subseteq \{1, 2, 3, 4\}$ and also $\{1, 2\} \subset \{1, 2, 3, 4\}$
- $\{1, 2, 3, 4\} \subseteq \{1, 2, 3, 4\}$ but it is not true that $\{1, 2, 3, 4\} \subset \{1, 2, 3, 4\}$
- set of men is a proper subset of the set of all people



Venn diagram:

Note: $A \subseteq A$, $\emptyset \subseteq A$ for every set A

Special Sets:

P - primes, N - natural numbers, Z - integers, $Q = \{\frac{a}{b}, a, b \in Z, b \neq 0\}$ - rational, R - real, I - irrational


$$P \subset N \subset Z \subset Q \subset R$$

Note: this simple theory of sets may lead to contradictions. Russell's paradox: Barber shaves only those men who do not shave themselves. Should the barber shave himself or not? Is the barber member of a set of men who shave themselves or not.

1.2 Basic Operations on Sets

There are ways to construct new sets from existing ones. Two sets can be "added" together, "subtracted", etc.

- **Union:** $A \cup B$ elements that belong to A or B .




Example: $\{1, 2\} \cup \{\text{blue, red}\} = \{1, 2, \text{blue, red}\}$

Properties:

- $A \cup B = B \cup A$
- $A \subseteq A \cup B$
- $A \cup A = A$
- $A \cup \emptyset = A$

- **Intersection:** $A \cap B$ elements that belong to A and B at the same time.



Example: $\{1, 2\} \cap \{\text{blue, red}\} = \emptyset$
 $\{1, 2\} \cap \{1, 2, 4, 7\} = \{2\}$


Properties:

- $A \cap B = B \cap A$
- $A \cap B \subseteq A$
- $A \cap A = A$

- $A \cap \emptyset = \emptyset$
- If $A \cap B = \emptyset$, then A and B are said to be disjoint.

• **Difference and Complement:** $B \setminus A$ or $B - A$: set of elements which belong to B , but not to A

In certain settings all sets under discussion are considered to be subsets of a given universal set U . Then, $U \setminus A$ is called complement of A and is denoted A' or A^C



Example: $\{1, 2, \text{green}\} \setminus \{\text{red}, \text{white}, \text{green}\} = \{1, 2\}$
 $\{1, 2\} \setminus \{1, 2\} = \emptyset$
 Integers \setminus Even numbers = Odd numbers

Properties:

- $A \cup A^C = U$
- $A \cap A^C = \emptyset$
- $(A^C)^C = A$
- $A \setminus A = \emptyset$
- $A \setminus B = A \cap B^C$

• **Cartesian product:** $A \times B$ combining every element from A with every element from B ; set of all ordered pairs (a, b) such that $a \in A$ and $b \in B$.


Example: $\{1, 2\} \times \{\text{red}, \text{white}, \text{blue}\} = \{(1, \text{red}), (1, \text{white}), (1, \text{blue}), (2, \text{red}), (2, \text{white}), (2, \text{blue})\}$

Properties:

- $A \times \emptyset = \emptyset$
- $A \times (B \cup C) = (A \times B) \cup (A \times C)$

Some identities:

- $A \setminus B = A \cap B^C$
- $(A \cup B)^C = A^C \cap B^C$
- $(A \cap B)^C = A^C \cup B^C$



Exercise 1: Do the following identities hold? You can find the answers to exercises at the end of this chapter.

(a) $(A \cap B) \cup (A \cap C) = A \cap (B \cup C)$

(b) $(A \cap B^c) \cup C = A \cap B$

Problem: 116 out of 129 students are regularly having lunch or supper. 62 students eat at most one of these meals. Further, there are 47 more students having lunch than students having supper. How many students are regularly having both lunch and supper? How many of them eat only supper and how many of them eat just lunch? [Hint: use Venn diagrams]

Solution: We will use the following notation:

x - students eating only lunch

y - students eating only supper

z - students eating both lunch and supper

w - students not eating any meal


Then the set-up of the problem says:

- $x + y + z = 116$
- $x + y + w = 62$
- $x = y + 47$

We know that $129 - 116$ students do not eat any meal, so $w = 129 - 116 = 13$.

Now we know that $x + y = 62 - 13 = 49$ and $x - y = 47$. This is true only for $x = 48$ and $y = 1$. Last, $x + y + z = 116$ implies that $z = 116 - 48 - 1 = 67$. Therefore, 67 students regularly eat both lunch and supper, 48 eat only lunch and 1 eats only supper.

1.3 Numbers



Real numbers (R): represented on real line with origin 0

Intervals: subsets of a real line

closed - e.g. $[2,5]$ - 2 and 5 belong to the interval

open - e.g. $(3,9)$ - 3 and 9 do not belong to the interval

Intersection: $[-4, 1] \cap [0, 2] = [0, 1]$

Union: $[-4, 1] \cup [0, 2] = [-4, 2]$



Example:

- $A = [-5, 3], B = [1, 10] \rightarrow A \cap B = [1, 3], A \cup B = [-5, 10]$
- $A = (-\infty, 2), B = (0, 4] \rightarrow A \cap B = (0, 2), A \cup B = (-\infty, 4]$
- $A = [-2, 8], B = (3, 10), C = (9, 15) \rightarrow A \cap B \cap C = \emptyset, A \cup B \cup C = [-2, 15)$

Exercise 2: Find intercept and union of the following intervals:

- $A = (0, 7], B = [1, 6]$
- $A = [-3, 4], B = (3, 10), C = (-1, 7)$

1.4 Answers

Exercise 1: The identity (a) holds and the identity (b) does not hold (to see this draw Venn diagrams).

Exercise 2:

- $A = (0, 7], B = [1, 6] \rightarrow A \cap B = [1, 6], A \cup B = (0, 7]$
- $A = [-3, 4], B = (3, 10), C = (-1, 7) \rightarrow A \cap B \cap C = (3, 4], A \cup B \cup C = [-3, 10)$

2 Logic

2.1 Simple Statement, Negation

A statement - a declarative sentence that is either true or false.

A simple statement - one that does not contain any other statement as a part.

Examples of sentences that are (or make) statements:

- "Socrates is a man."
- "A triangle has three sides."
- "Paris is the capital of England."

The first two (make statements that) are true, the third is (or makes a statement that is) false.

Examples of sentences that are not (or do not make) statements:

- "Who are you?"
- "Run!"

Negation - In logic and mathematics, negation or "not" is an operation on logical values, which changes true to false and false to true. Intuitively, the negation of a proposition holds exactly when that proposition does not hold.

Notation: statement - p , can be true or false; if p is true then "NOT p " or " $\sim p$ " or " $\neg p$ " is false.

Examples of negations of previous statements:

- "Socrates is not a man." or "Socrates is a woman."
- "A triangle does not have three sides." or "A triangle has at most two or at least four sides."
- "Paris is not the capital of England."

Notation: \forall - for all/every/each; \exists - there exists/at least one

Examples: A is a set of all natural numbers larger than 10: $A = \{\forall x \in N; x > 10\}$

There exists at least one number for which the square root is negative: $\exists x; \sqrt{x} < 0$

Examples: Decide if the following statements are true or false and find negations:

- Math teacher is nice. True. Math teacher is not nice.
- There are **no** unemployed people in the CR. False. There is **at least** one unemployed person in the CR.
- **At least four** students in class are women. True (most of the time). **At most three/less than four** students in class are women.
- Today is Tuesday. False (depends on when you read this material). Today is not Tuesday.

Exercise 1: Decide if the following statements are true or false and find negations:

- **At most three** students have blue eyes.
- Every day has 25 hours.
- **All** cars are red.
- The semester begins today.

2.2 A Compound Statement

A compound statement - one with two or more simple statements as parts

An operator - joins simple statements into compounds

Compound statements:

statement	symbol	how we read it
Conjunction	$p \wedge q$	both p and q are true
Disjunction	$p \vee q$	either p or q is true or both
Implication	$p \Rightarrow q$	if p is true then q is true
Equivalence	$p \Leftrightarrow q$	p and q are either both true or both false

Examples:

- Conjunction: I had an ice-cream *and* a cake.
Tom is tall *and* has dark hair.
- Disjunction: I had a coffee *or* tea.
I went to cinema *or* for a dinner.
- Implication: *If* you study hard (*then*) you will get a good grade.
If it rains (*then*) I will stay at home.
- Equivalence: I will go to cinema *if and only if* (*iff*) you go.
The flag is on the castle *if and only if* (*iff*) the president is inside.